

Charges and Currents in the Noncommutative Chern-Simons Theory of the QHE

T. H. Hansson and A. Karlhede

*Department of Physics, University of Stockholm,
Stockholm Center for Physics, Astronomy and Biotechnology
S-11385 Stockholm, Sweden*

(February 1, 2008)

We couple the noncommutative Chern-Simons theory describing the fractional quantum Hall effect to external magnetic and electric potentials, and derive expressions for charge and current densities. To lowest non-trivial order the density operator satisfies the commutator algebra characteristic of the lowest Landau level. We take this as a strong indication that the theory correctly describes point particles in a strong magnetic field. We also obtain the correct QH response to a constant electric field, and density modulations in response to a weak variable magnetic field.

PACS numbers: 73.43.-f, 11.15.-q

It is well established that the low-energy effective lagrangian for a quantum Hall system is a Chern-Simons (CS) gauge theory. Specifically, for the simplest case of the Laughlin fractions, it takes the form,

$$\mathcal{L} = \frac{k}{4\pi} \epsilon^{\mu\nu\lambda} a_\mu \partial_\nu a_\lambda - \frac{e}{2\pi} \epsilon^{\mu\nu\lambda} A_\mu \partial_\nu a_\lambda, \quad (1)$$

where k is an odd integer, and A_μ is the external electromagnetic potential. This lagrangian (and its generalizations to other fractions, multi-component systems *etc.*) correctly describes the quantum Hall conductance, and also ground state degeneracies on higher genus surfaces and edge excitations [1]. At the level of the effective theory, the quantization of k does not follow from any general principle but should be regarded as an input. There are, however, microscopic derivations of (1), relying on various mean field approximations, where the quantization of k essentially follows from a statistical transmutation [2].

Another limitation of (1) follows upon the realization that the physics relevant for the Laughlin fractions of the QHE can be understood entirely in terms of states in the lowest Landau level (LLL). In particular, this means that the Fourier components of the density operator do not commute, but obey

$$[\rho_{\vec{k}}, \rho_{\vec{p}}] = -2i \sin\left(\frac{\ell^2}{2}(\vec{k} \times \vec{p})\right) e^{\frac{\ell^2}{2}\vec{k} \cdot \vec{p}} \rho_{\vec{k}+\vec{p}}, \quad (2)$$

where $\ell = (\hbar/eB)^{1/2}$ is the magnetic length [3]. In the effective theory (1), however, $\rho(x) = 2\pi\epsilon^{ij}\partial_i a_j = 2\pi b$, and the commutator (2) is $\sim (\vec{k} \times \vec{p})\delta(\vec{k} + \vec{p})$.

Recently, Susskind proposed a noncommutative version of (1),

$$\mathcal{L}_{nc} = \frac{k}{4\pi} \epsilon^{\mu\nu\lambda} \left(\hat{a}_\mu \star \partial_\nu \hat{a}_\lambda + \frac{2i}{3} \hat{a}_\mu \star \hat{a}_\nu \star \hat{a}_\lambda \right), \quad (3)$$

where the noncommutative parameter entering the Moyal star product is $\theta = 1/2\pi\rho$ with ρ being a constant background density [4]. Susskind argues that θ can be interpreted as the unit area per particle, and that (3) incorporates the particle substructure into the effective theory. That this is an essential point when constructing effective actions was emphasized by Haldane in the case of one dimensional quantum fluids [5]. For further discussion of the noncommutative CS (NCCS) theory we refer to [4,6–10]. Here we just mention that invariance under large gauge transformations requires k to be an integer independent of the physical interpretation of the NCCS action [11,6], while the restriction to odd integers is a consequence of the electrons forming the QH fluid being fermions [4].

In this letter we extend the NCCS theory to include coupling to an external electromagnetic field. We then derive expressions for charge and current densities, verify the commutation relation (2) to $O(\hbar\theta^0)$ in an expansion in $\hbar$ and θ . We also show that a weak variable magnetic field gives rise to density fluctuations, and that the response to a constant electric field is given by the Hall conductance $\sigma_{ab} = \sigma_H \epsilon_{ab}$. These results lend further support to the notion that the NCCS theory is the correct low energy theory of the QHE.

There is an equivalent formulation of NCCS theory in terms of a matrix model [12]. To include couplings to external potentials, we propose the following extension of the matrix version of Susskind's lagrangian,

$$L = \frac{eB}{2} \text{Tr} \left\{ \left(\dot{X}^a - i[X^a, \hat{a}_0]_m \right) \left(\epsilon_{ab} X^b + \frac{2}{eB} \hat{A}_a \right) + 2\theta \hat{a}_0 - \frac{2}{eB} \hat{A}_0 \right\}, \quad (4)$$

where $X^a(t)$ are hermitian matrices, and $\hat{a}_0(t)$ is a matrix Lagrange multiplier imposing the (classical) commutation relation

$$[X^a, \epsilon_{ab} X^b + \frac{2}{eB} \hat{A}_a]_m = 2i\theta. \quad (5)$$

The new terms are the couplings to the matrices, $\hat{A}_\mu(X^a, t)$ ($\mu = 0, 1, 2$; $a, b = 1, 2$), which are related to the ordinary gauge potential, $A_\mu(x^a, t)$, by the Weyl ordering formula,

$$\hat{f}(X^a, t) = \int \frac{d^2 k d^2 x}{(2\pi)^2} f(x^a, t) e^{ik_a(x-X)^a}, \quad (6)$$

where $f(x^a, t)$ is an ordinary function of the usual (commuting) coordinate x^a . To exhibit the canonical structure of the theory, we choose a linear gauge, $\hat{A}_2 = 0$ and rewrite the constraint (5) as,

$$[X^1, X^2 + \frac{1}{B} \hat{A}_1]_m = [X^1, \frac{1}{B} P^1]_m = i\theta, \quad (7)$$

where P^1 is the momentum conjugate to X^1 as calculated from the lagrangian (4). For $\hat{A}$ corresponding to a constant external magnetic field, we get $[X^1, X^2] = \text{const.}$

In the $\hat{a}_0 = 0$ gauge, the lagrangian (4) is quantized by imposing equal time (quantum) commutation relations,

$$[X_{mn}^1, P_{rs}^1] = i\hbar \delta_{ms} \delta_{nr}. \quad (8)$$

Note that expressed in canonical variables, the constraint takes the same form as in the $A_\mu = 0$ case. Thus the whole discussion about constraints, exchange phases etc. will be the same as in refs. [4] and [6].

We now proceed to analyze the proposed lagrangian (4). The gauge transformation $\delta A_\mu(x) = \partial_\mu \Lambda(x)$ implies the following transformation for the corresponding matrices,

$$\begin{aligned} \delta \hat{A}_{mn}^a &= - \left(\frac{\partial \hat{\Lambda}}{\partial X^a} \right)_{mn} = - \frac{\partial \text{Tr} \hat{\Lambda}}{\partial X_{nm}^a} \\ \delta \hat{A}_0 &= \partial_0 \hat{\Lambda}, \end{aligned} \quad (9)$$

where again the matrix valued function $\hat{\Lambda}(X^a, t)$ is related to $\Lambda(x^a, t)$ by the Weyl ordering (6), and the derivative satisfies $\partial e^{-ik_b X^b} / \partial X^a = -ik_a e^{-ik_b X^b}$. It is straightforward to show that the lagrangian (4) is invariant both under the ‘background’ gauge transformation (9), and the noncommutative gauge transformation,

$$\begin{aligned} \delta X^a &= i[\hat{\Lambda}, X^a]_m \\ \delta \hat{a}_0 &= i[\hat{\Lambda}, \hat{a}_0]_m + \partial_0 \hat{\Lambda}. \end{aligned} \quad (10)$$

Inserting the proper Weyl ordered expression for $\hat{A}$ into (4) and taking the variation of the corresponding action *w.r.t.* A_μ , we obtain the following expressions for the Fourier components of the particle and current densities,

$$\begin{aligned} \rho_{\vec{k}} &= \text{Tr} \left(e^{-ik_a X^a} \right) \\ j_{\vec{k}}^a &= \text{Tr} \left(\dot{X}^a e^{-ik_b X^b} \right), \end{aligned} \quad (11)$$

which clearly are invariant under both (9) and (10). Note that the simple commutation relation (7) is between X^1 and $P^1 = eBX^2 + \hat{A}_1$, not between X^1 and X^2 - this makes it hard to evaluate the densities (11).

A noncommutative gauge theory does not allow for local observables, and the expressions (11) are in fact closely related to the so called open Wilson loops [13]. Note that $\rho_{\vec{k}}$ is the natural noncommutative generalization of the usual expression

$$\tilde{\rho}_{\vec{k}} = \int d^2 r e^{-i\vec{k} \cdot \vec{r}} \rho(\vec{r}) = \int d^2 r e^{-i\vec{k} \cdot \vec{r}} \sum_{i=1}^N \delta^2(\vec{r} - \vec{r}_i) = \sum_{i=1}^N e^{-i\vec{k} \cdot \vec{r}_i}. \quad (12)$$

The connection between the commutative and noncommutative case can be made more precise in the modified theory due to Polychronakos, where the constraint (5) is relaxed in a minimal way in order to allow for a finite system

of N particles described by $N \times N$ matrices [6]. Up to finite size corrections, which should vanish in the large N limit, we still expect the current and charge densities to be given by (11). In the finite N formulation one can pick a gauge where the $2N$ physical degrees of freedom are the diagonal elements of X^a . One of the matrices, X^1 say, can be diagonalized, while the off diagonal elements of the other are given by $X_{mn}^2 = i\theta/(x_m - x_n)$, where x_n are the diagonal elements of X^1 . In the limit of large x^a and large separation compared to $\sqrt{\theta}$, both matrices are almost diagonal and the Fourier transform of $\rho_{\vec{k}}$ in (11) reproduces $\rho(\vec{r})$ in (12) with $\vec{r}_n = (x_n, y_n)$. This confirms the interpretation of the diagonal elements of X^a as the positions of the electrons.

The densities (11) are not local, and one would hope that this non-locality would reflect the LLL physics of the QH effect. We can now test this proposition by evaluating the commutator (2) using our density operator (11). This calculation is cumbersome, since it involves commutators of exponentials of matrices where the elements do not commute. However, using series expansions it is relatively straightforward to obtain

$$[\rho_{\vec{k}}, \rho_{\vec{p}}] = -i\ell^2(\vec{k} \times \vec{p})\rho_{\vec{k}+\vec{p}} + O(\hbar\theta, \hbar^2), \quad (13)$$

which we recognize as the leading term in an $\hbar$ expansion of (2). (The sign is related to the sign of B via (8), and we follow the convention of refs. [4] and [6].)

Note that already to this order the (classical) matrix structure must be taken into account in order to reproduce the $\rho_{\vec{k}+\vec{p}}$ term characteristic for the LLL projection, but also note that the result is insensitive to the ordering of the exponential in (11), so to test the Weyl ordering prescription one must go to higher orders in θ . The lagrangian (4), the particle and current density operators (11), and the commutator (13) are the main results of this paper.

It is in general hard to calculate the response to an external magnetic field since (6) only gives an implicit relation for the matrix $\hat{A}^a(X^1, P^1)$. It can however be done for a weak perturbing field of the form $\delta B(\vec{x}) = \epsilon B \sin(\vec{q} \cdot \vec{x})$, where $\epsilon \ll 1$. Substituting this in (6) and (11), and expanding to lowest order in ϵ , the trace can be evaluated to lowest order in θ using a coherent state basis, to yield

$$\rho(\vec{x}) = \frac{1}{2\pi\theta} [1 + \epsilon \sin(\vec{q} \cdot \vec{x})] + O(\theta^0) \quad (14)$$

as expected.

Next we calculate the current response to an applied constant electric field of the form, $A_0 = E_a x^a$. To evaluate (11), we first need to determine $\dot{X}^a$ from the equations of motion, a problem very similar to the QH droplet in a quadratic matrix potential considered by Polychronakos [6]. Obtaining $\dot{A}_0$ from (6), substituting in (4) and varying with respect to X^a , yields

$$B\dot{X}^a = -\epsilon_{ab}\hat{E}^b. \quad (15)$$

Substituting this into the expression (11) gives the following expression for the electromagnetic current density $J^a = ej^a$,

$$J_k^a = -\epsilon_{ab} \frac{eE^b}{B} \rho_{\vec{k}}, \quad (16)$$

and taking the Fourier transform we finally get

$$J^a = -\nu \frac{e^2}{h} \epsilon_{ab} E^b = -\sigma_H \epsilon_{ab} E^b, \quad (17)$$

where the filling fraction ν is defined by $\rho = \nu eB/h$. From ref. [4] we know that for the particles described by the matrix model to be fermions, we must have $\nu^{-1} = 2n + 1$ for which (17) gives the Laughlin values for the quantized Hall conductance σ_H .

To conclude, we have presented substantial reasons to believe that the noncommutative Chern-Simons Lagrangian (4) correctly describes the coupling of a Laughlin QH liquid to external electromagnetic potentials. That the density commutator (13), characteristic of the lowest Landau level, emerges from our construction provides further evidence that the theory in fact describes point particles in a strong magnetic field.

It is an obvious and interesting problem to evaluate the commutator (13), and the density response (14) to higher order in $\hbar$ and θ . Such calculations should certainly be done in order to see if the full $\vec{k}$ and $\vec{p}$ dependence in (2) is reproduced by the operators (11). This will show whether the full LLL physics is hidden in (4), or only the leading long distance part of it. That there exists a mapping from the NCCS theory to the Calogero-Sutherland model [6],

and also a connection to the Laughlin wavefunctions [9], suggest the former scenario, although the mapping might be quite complicated [10]. Finally it should be straightforward to modify our construction to the finite droplet considered in ref. [6]. This would hopefully allow for a more detailed analysis of edge modes and edge currents.

Acknowledgement: We thank Alexios Polychronakos for pointing out errors in the first version of this paper, and for helping us to get the correct form of eq. (9).

-
- [1] For a review, see *e.g.* A. Zee, “*Quantum Hall fluids*” in Proc. of South African School of Physics, Tsitsikamma - 94; Springer Verlag. [cond-mat/9501022].
 - [2] See *e.g.* the review S. C. Zhang, Int. J. Mod. Phys. **6**, 25 (1992).
 - [3] S. M. Girvin, A. H. MacDonald, and P. M. Platzman, Phys. Rev. B **33**, 2481 (1986).
 - [4] L. Susskind, hep-th/0101029.
 - [5] F. D. M. Haldane, Phys. Rev. Lett. **47**, 1840 (1981).
 - [6] A. P. Polychronakos, JHEP **0104**, 011 (2001) [hep-th/0103013].
 - [7] A. P. Polychronakos, hep-th/0106011;
 - [8] B. Morariu and A. P. Polychronakos, hep-th/0106072.
 - [9] S. Hellerman and M. Van Raamsdonk, hep-th/0103179.
 - [10] D. Karabali and B. Sakita, hep-th/0106016.
 - [11] V. P. Nair and A. P. Polychronakos, hep-th/0102181.
 - [12] A. P. Polychronakos, JHEP **0011**, 008 (2000) [hep-th/0010264].
 - [13] For a review, see M. R. Douglas and N. A. Nekrasov, hep-th/0106048.